

SHORT NOTE

Remote Sensing and Depth Distribution of Ocean Chlorophyll

Raymond C. Smith

Visibility Laboratory, Scripps Institution of Oceanography, University of California, San Diego, La Jolla, California 92093, USA

ABSTRACT: This note considers how chlorophyll, as estimated from Coastal Zone Color Scanner (Nimbus 7 satellite) data, is related to the chlorophyll concentration in the oceanic water column. In spite of some limitations, the remotely sensed chlorophyll concentration can be used as an index of the mean water column chlorophyll. Chlorophyll imagery provides synoptic measurements, impractical to obtain from ships alone, and hence may be of considerable importance in phytoplankton ecology.

It has been demonstrated that data from the Coastal Zone Color Scanner (CZCS) on the Nimbus 7 satellite can be processed to provide quantitative chlorophyll maps of an oceanographic region within an accuracy of $0.5 \log C$ (Morel and Prieur, 1977; Gordon et al., 1980; Hovis et al., 1980; Smith and Baker, submitted, 1981). This note discusses the issue of how chlorophyll, as estimated from CZCS imagery, is related to the water column chlorophyll concentration.

The problem is best explored in terms of optical attenuation lengths as distinct from geometrical depths. Gordon and McCluney (1975) defined, Z_{90} , as the depth of penetration of light above which 90 % of the diffusely reflected irradiance (excluding specular reflectance) originates. They showed that for a homogeneous ocean

$$Z_{90} \approx K^{-1} \quad (1)$$

where K = diffuse attenuation coefficient for downwelling irradiance. More recently, it has been shown (Gordon and Clark, 1980), that for remote sensing purposes, the concentration of the constituent under consideration should be weighted by a factor

$$g(Z) = \exp \left\{ -2 \int_0^Z K(Z) \cdot dZ \right\} \quad (2)$$

Frequently, as a first-order approximation, $K(Z)$ is considered approximately constant with depth so that,

$$g(Z) = \exp \left\{ -2K \cdot Z \right\} \quad (3)$$

Heuristically, this weighting factor can be viewed as being derived from the irradiance arriving at the surface having been attenuated by $\exp[-K \cdot Z]$ from the surface to the depth Z and by the same factor on the return to the surface. Gordon and Clark (1980) conclude that the remotely sensed concentration of chlorophyll is given by

$$C_{SAT} = \frac{\int_0^{Z_{90}} C(Z) \cdot g(Z) \cdot dZ}{\int_0^{Z_{90}} g(Z) \cdot dZ} \quad (4)$$

where $C(Z)$ is the concentration of chlorophyll as a function of depth.

When shipboard chlorophyll has been determined at only a few discrete depths in the water column, then C_{SAT} may be approximated by the mean chlorophyll concentration to a depth of one attenuation length, C_K (Smith and Baker, 1978a). The mean chlorophyll to the euphotic depth, C_T , is obtained from an integral of $C(Z)$ over 4.6 attenuation lengths (i.e. the 1 % level). If, as is generally the case, $C(Z)$ is not too complex then C_{SAT} , C_K and C_T are highly correlated, as has been previously shown using data from 140 stations (Smith and Baker, 1978a). Thus C_{SAT} can be used as an index of the mean chlorophyll concentration in the water column.

In order to illustrate this for a variety of water types, Fig. 1 and Table 1 show 3 chlorophyll profiles (Cullen and Eppley, 1981) ranging from productive coastal water to oligotrophic waters. It is useful to renormalize these profiles as shown in Fig. 2 where chlorophyll has been plotted as a function of attenuation lengths of photosynthetic available radiant energy (PAR). When plotted in this manner, a given percentage PAR level (the 1 % euphotic depth for example) and/or the




Fig. 1 Typical chlorophyll profiles plotted as function of geometrical depth (Cullen and Eppley, 1980): (A) North Pacific Central Gyre, near 28° N, 155° W (from Beers et al., 1975); (B) SCBS 15, Sta 205; (C) SCBS 7, Sta. 102

remote sensing penetration depth are directly comparable for waters with very different chlorophyll concentrations. Thus, C_{SAT} for these very different profiles would be proportional to the integral of the weighting function, $g(Z)$, times the profiles as shown in Fig. 2 and as given by Eq. (4). Fig. 2 illustrates that C_{SAT} is determined by a weighting of the top 20 to 30 % of the euphotic zone, regardless of whether oligotrophic or highly productive conditions exist.

Several observations with respect to Table 1 can be made. First, as one moves from oligotrophic to eutrophic waters the euphotic depth and the PAR penetration depth change together proportionally. Second, the

Table 1 Observations on 3 typical vertical profiles (after Cullen and Eppley, 1981): (A) North Pacific Central Gyre (Beers et al., 1975) and Food Chain Research Group (Eppley, unpubl.); (B) Offshore Southern California Bight, SCBS 15, Sta. 205; (C) Nearshore Southern California Bight, SCBS 7, Sta. 102

	A	B	C
Euphotic depth (m)	105	66	33
PAR penetration depth			
K_{PAR}^{-1} (m)	22.8	14.3	7.2
Integrated Chl ($mg\ m^{-2}$)	18.5	19.8	20.2
Mean Chl a , C_T ($mg\ m^{-3}$)	0.09	0.38	0.67
Satellite estimated Chl			
C_{SAT} ($mg\ m^{-3}$)	0.045	0.13	0.36
Integrated primary production			
P_I ($gm\ m^{-2}\ d^{-1}$)	0.054	0.162	0.516




Fig. 2. Same profiles as in Fig. 1 plotted as function of attenuation lengths of PAR ($K_{PAR} \cdot Z$). The remote sensing weighting function, $g(Z)$, is shown as dashed curve. Depth of euphotic zone, taken as 1% PAR level, is indicated by an arrow on vertical scale

mean chlorophyll concentrations in the water column (to the 1 % level), C_T , varies in a manner similar to C_{SAT} . Third, both C_{SAT} and C_T are correlated with the integrated primary production in the water column.

The wavelength dependence of K is also important when considering the penetration depth. For calculations involving primary productivity it is generally most useful to consider a broadband width K as the attenuation coefficient for PAR, defined so that

$$PAR(Z) = PAR(O^-) \exp(-K_{PAR} \cdot Z) \quad (5)$$

where K_{PAR} is the diffuse attenuation coefficient for PAR (350 to 700 nm). However, K is also a function of each narrow band wavelength, in particular the four visible CZCS spectral bands (443, 520, 550, and 670 nm). The CZCS spectral bands are relatively narrow ($\Delta\lambda = 20$ nm FWHM) so the depth of penetration will vary with each band. Our bio-optical model (Smith and Baker, 1978b; Baker and Smith, submitted, 1981), which gives the relationship between $K(\lambda)$ and C , can be used to show how the penetration depth varies with chlorophyll concentration in each CZCS spectral band.

In Fig. 3 the penetration depth, Z_{90} , has been calculated for four CZCS bands and for K_{PAR} as a function of chlorophyll concentration. This figure illustrates several points related to the remote sensing of chlorophyll. First, the penetration depth for all spectral bands becomes smaller as the chlorophyll concentration increases. Second, the penetration depth for PAR and the 520 nm band are nearly the same regardless of variations in C . This is consistent with the concept of a

'hinge point' (Duntley et al., 1974), which is a wavelength band where changes in the upwelled spectral radiance are relatively insensitive to variations in chlorophyll. Third, K_{070}^{-1} is only a few meters and relatively independent of C because the attenuation in the red is dominated by the absorption of water. Fourth, the penetration depth at 550 and 443 nm generally differ from K_{PAR}^{-1} . Indeed, it is the difference in these bands that makes the remote sensing of chlorophyll possible.

There are a number of reasons why a remotely sensed estimate, C_{SAT} , of the mean water column chlorophyll concentration, C_T , may be inexact: (1) normalization to single broadband penetration depth, K_{PAR}^{-1} , while a good first approximation, neglects the wavelength dependency of the band width penetration depth used in the chlorophyll algorithm; (2) significant




Fig. 3. Penetration depth, $Z_{(H)} \approx K^{-1}$ (m), versus chlorophyll concentration, C ($\text{mg} \cdot \text{m}^{-3}$) as derived from an optical classification model (Smith and Baker, 1978b; Baker and Smith, 1981). Penetration depth calculated for 4 CZCS spectral bands K_{λ}^{-1} and for total photosynthetically available radiant energy, K_{PAR}^{-1} . Scale on right ordinate gives ratio of penetration depth in blue band to that in the green, $K_{443}^{-1}/K_{550}^{-1}$.

changes in the chlorophyll profiles at depth greater than a few attenuation lengths, without a proportional change in the top region of the profile, will go unnoticed by the satellite sensor; (3) shifts in the ratio of chlorophyll-like pigments (e.g. chlorophyll a and phaeophytin a) with depth do not change the optical signal.

In spite of these limitations, the examples illustrate that C_{SAT} can be used as an index of the mean water column chlorophyll. The accuracy of this index has been studied and shown to be within less than 0.5 log C (Gordon et al., 1980; Smith and Baker, submitted, 1981). Since chlorophyll imagery can provide synoptic measurements impractical to obtain from ships alone, the CZCS data can be of considerable value in studies of phytoplankton ecology especially in complex coastal regions.

Acknowledgements. This work was completed with the support of NASA Grant No. NSG 1641. K. S. Baker and R. W. Eppley provided valuable discussions and advice.

LITERATURE CITED

- Baker, K. S., Smith, R. C. (1981). Bio-optical classification and model of natural waters II. (submitted)
- Beers, J. R., Reid, F. M. H., Stewart, G. L. (1975). Microplankton of the North-Pacific Central gyre. Population structures and abundance, June 1973. *Int. Revue ges. Hydrobiol.* 60: 607-638
- Cullen, J. J., Eppley, R. W. (1981). Chlorophyll maximum layers of the Southern California Bight and possible mechanisms of their formation and maintenance. *Oceanol. Acta* 4: 23-32
- Duntley, S. Q., Austin, R. W., Wilson, W. H., Edgerton, C. F., Moran, S. E. (1974). Ocean color analysis. *Scripta Inst. Oceanogr. Ref.* 74-10
- Gordon, H. R., Clark, D. K. (1980). Remote Sensing optical properties of a stratified ocean. *Appl. Opt.* 19: 3428-3430
- Gordon, H. R., Clark, D. K., Mueller, J. L., Hovis, W. A. (1980). Phytoplankton pigments from the Nimbus-7 Coastal Zone Color Scanner: comparison with surface measurements. *Science, N.Y.* 210: 63-66
- Gordon, H. R., McCluney, W. R. (1975). Estimation of the depth of sunlight penetration in the sea for remote sensing. *Appl. Opt.* 14: 413-416
- Hovis, W. A., Clark, D. K., Anderson, F., Austin, R. W., Wilson, W. H., Bkaer, E. T., Ball, D., Gordon, H. R., Mueller, J. L., El-Sayed, S. Z., Sturm, B., Wrigley, R. C., Yentsch, C. S. (1980). Nimbus-7 Coastal Zone Color Scanner Systems: description and initial imagery. *Science, N.Y.* 210: 60-63
- Morel, A., Prieur, L. (1977). Analysis of variations in ocean color. *Limnol. Oceanogr.* 22: 709-722
- Smith, R. C., Baker, K. S. (1981). Oceanic Chlorophyll concentrations as determined using Coastal Zone Color Scanner imagery. *Mar. Biol.* (submitted)
- Smith, R. C., Baker, K. S. (1978a). The bio-optical state of ocean waters and remote sensing. *Limnol. Oceanogr.* 23: 247-259
- Smith, R. C., Baker, K. S. (1978b). Optical classification of natural waters. *Limnol. Oceanogr.* 23: 260-267

Accepted for printing on May 15, 1981