

Spectral beam attenuation coefficient retrieved from ocean color inversion

Collin S. Roesler

Bigelow Laboratory for Ocean Sciences

Emmanuel Boss

School of Marine Sciences, University of Maine

Abstract. A reflectance inversion model is presented in which particle backscattering is parameterized by the particle backscattering ratio, attenuation and absorption coefficients rather than a hyperbolic function. This parameterization removes backscattering spectral constraints and yields independent estimates of the magnitude and spectral slope of the particle beam attenuation, which have been found to correlate with the particle concentration and size distribution, respectively. The model yields estimates of component absorption and backscattering consistent with observations and Mie models, particularly for environments dominated by phytoplankton and other strongly absorbing particles.

1. Introduction

Ocean color contains a wealth of information regarding the concentration and composition of dissolved and suspended constituents in the sea. Analytic expressions derived from simplifications to the radiative transfer equation express the radiance and irradiance reflectance as a function of the inherent optical properties (IOPs) [e.g., *Gordon et al.*, 1988]:

$$R = G \frac{b_b}{a + b_b} \quad (1)$$

where R is reflectance (ratio of either upward irradiance or radiance to the downward irradiance), G depends upon the definition of R and water and atmospheric properties, and a and b_b are absorption and backscattering coefficients.

A number of semi-analytical inversion models extract constituent IOPs from measured reflectance signatures [*Gordon et al.*, 1988; *Lee et al.*, 1994; *Roesler and Perry*, 1995; *Hoge and Lyon*, 1996; 1999; *Garver and Siegel*, 1997]. The IOPs are linearly deconvolved into operational components and expressed as the product of scalar amplitudes and non-dimensionalized basis functions (spectra):

$$a = a_w + L_\varphi \hat{a}_\varphi + L_{\text{CPM}} \hat{a}_{\text{CPM}} + L_{\text{CDM}} \hat{a}_{\text{CDM}} \quad (2a)$$

$$b_b = b_{bw} + L_{\text{bbp}} \hat{b}_{\text{bp}} \quad (2b)$$

where the subscripts w , p , φ , CPM and CDM indicate water, total particles, phytoplankton, colored particulate material (exclusive of phytoplankton), and colored dissolved material, amplitudes are indicated by L , and basis functions by $\hat{\cdot}$.

Published inversion models differ in basis function definitions and solution methods. Decades of laboratory and field investigations yield robust constraints on the absorption basis function spectral variations. Backscattering sensors are relatively new and there is less information on backscattering

basis functions. Both observations and Mie theory for absorbing spheres demonstrate strong spectral features near absorption peaks [*van de Hulst*, 1957; *Gordon*, 1974; *Bricaud and Morel*, 1986; *Zaneveld and Kitchen*, 1995]. Difficulty in constraining these features, has lead to implementation of Mie theory for populations of *non-absorbing* homogeneous spheres, in which backscattering is expressed as a smoothly varying function [*Morel*, 1973]:

$$b_{bp}(\lambda) = b_{bp}(\lambda_o) \left(\frac{\lambda}{\lambda_o}\right)^\eta \quad (3)$$

where λ is wavelength, λ_o is a nominal scaling wavelength, and η is the hyperbolic slope (which is either held constant or retrieved by inversion).

The utility of such models has been demonstrated over a range of environments, particularly for estimation of a_p and the combined $a_{CPM+CDM}$, with less confidence in derived b_{bp} values. In this paper, we propose to build upon this background to develop an algorithm that removes spectral constraints on b_{bp} by allowing the influence of a_p on b_{bp} to be parameterized. This is achieved by using IOP observations and theory to expand the inversion algorithm and in doing so retrieve estimates of the spectral particle beam attenuation coefficient and the particle backscattering ratio.

2. Model Development and Tests

The focus of this model relies on replacing the expression for b_{bp} given by (3), in which we have relatively little confidence, by a function in whose spectral shape we have a good deal of confidence. We express the particle backscattering spectrum as:

$$b_{bp}(\lambda) = \tilde{b}_{bp}(\lambda) b_p(\lambda) \quad (4)$$

where \tilde{b}_{bp} is the particle backscattering ratio (defined as the backscattering to total scattering ratio), and b_p is the particle scattering spectrum. Mie theory predicts spectrally independent backscattering ratios for Junge populations of absorbing particles with constant real refractive indices [*Ulloa et al.*, 1994]. Although some spectral variations are observed for monospecific phytoplankton cultures [*Stramski and Mobley*, 1997], in situ observations on limited data sets suggest they are weak [$<10\%$, *Twardowski et al.*, 2001]. Assuming no spectral dependence in the backscattering ratio and recognizing that b_p can be expressed by the difference of a_p from c_p yields:

$$b_{bp}(\lambda) = \tilde{b}_{bp} [c_p(\lambda) - (a_p(\lambda) + a_{CDM}(\lambda))] \quad (5)$$

Unlike the spectral dependence of b_{bp} , c_p is observed to be a smoothly varying function of wavelength, even for strongly absorbing particle populations [*Voss*, 1992; *Barnard et al.*, 1998; *Boss et al.*, 2001a], and is well described by:

$$c_p(\lambda) = c_p(\lambda_o) \left(\frac{\lambda}{\lambda_o}\right)^\gamma \quad (6)$$

where γ is the hyperbolic slope. Mie solutions for c_p show small departures from (6) for hyperbolically distributed absorbing particles ($<2\%$) [*Boss et al.*, 2001b]. We define the c-model for inversion (c as in attenuation) by (1) with F/Q replace by the estimable amplitude $L_{F/Q}$, a given by (2a) and:

$$b_b = b_{bw} + L_{bbp} \left[L_{cp} \left(\frac{\lambda}{\lambda_0} \right)^{-c_\gamma} - L_\phi \hat{a}_\phi - L_{CPM} \hat{a}_{CPM} \right] \quad (7)$$

The basis functions are defined *a priori* and amplitudes, L , are estimated by optimization. The spectral restriction on b_{bp} , as defined by (3), is replaced by two functions, \tilde{b}_{bp} and c_p , which have minimal or well-constrained spectral dependences.

We solve the inversion with a bounded non-linear least squares approach (Levenberg-Marquard; [Press *et al.*, 1989]). Without bounds, >1 minima can be found over a three-order dynamic range yielding negative/non-physical solutions. We define *a priori* three overlapping ranges of upper and lower amplitude bounds (where each range spans over a factor of 10) based upon environment: open ocean, coastal and extreme blooms. For example, the amplitude bounds for L_{cp} are 0.03-0.3, 0.1-1, and 0.8-8.

The amplitude $L_{F/Q}$ is allowed to vary within the entire F/Q range observed by Morel and Gentili [1996], but not spectrally, although it has been shown to have spectral variations [Morel *et al.*, 2002]. As we gain more understanding of the spectral variations, this factor can become a basis function in the model. Water backscattering [Morel, 1974] and absorption [Pope and Frye, 1997] are corrected for in situ temperature and salinity conditions [Morel, 1974; Pegau *et al.*, 1997]. The basis functions for \hat{a}_{CDM} and \hat{a}_{CPM} are defined by exponential functions with slopes of -0.018 and -0.011 nm^{-1} , respectively [Roesler *et al.*, 1989]. For simplicity, the basis function for phytoplankton, \hat{a}_ϕ , is defined by an average spectrum (Fig. 1 in Roesler and Perry [1995]), although using multiple functions for each pigment-based taxonomic groups improves reflectance fit and yields taxonomic information [Roesler *et al.* 2003].

C-model results are compared first to those derived from the published model of Roesler and Perry [1995] (referred to here as the *standard model*). Identical basis functions were used for the \hat{a} components in both models. In the standard model b_{bp} was parameterized, as published, by the sum of 2 basis functions defined by (3) with $\eta=0$ and -1 , respectively.

Amplitudes from the c-model were then compared with measured values collected over a broad range of optical domains: clear gyre waters off the Oregon coast, coastal Gulf of Maine (GoM) and Benguelan Upwelling (BU) waters, some with extreme algal bloom conditions. Observed a_p , a_{CDM} , and c_p spectra were derived from in situ ac_9 measurements using simultaneous filtered (0.2- μm) and unfiltered configurations, with corrections as in Roesler [1998]. The spectral slope, γ , was derived by non-linear least-squares regression of (6) onto c_p observations. b_{bp} coefficients were measured with a WETLabs ECOvsf, calibrated with beads and corrected for variable path absorption. There were no backscattering measurements from the gyre and GoM stations, so b_{bp} and \tilde{b}_{bp} were estimated from Petzold [1972] using data from waters off Bermuda (AUTEC) and offshore southern California (HAOCE), respectively, which had absorption and attenuation coefficients within 10% of our measured values.

Hyperspectral surface reflectance was measured with either a LICOR 1800UW scanning radiometer [Roesler and Perry, 1995] or a Satlantic H-TSRB radiometer buoy. Upwelling

measurements were corrected to the surface using spectral attenuation coefficients measured in the top meter.

3. Results

Chlorophyll values ranged from 0.07 to 131.65 $\mu\text{g/L}$. Reflectance spectra were well predicted by both models although the c-model yielded better fits (Fig. 1A, B). The spectral shape of b_{bp} was better predicted by the c-model, with the standard model often yielding unrealistic spectra (e.g. negative values, Fig. 1C, D). The c-model provided significantly better estimates of a_p and a_{CDM} amplitude and spectral shape (Fig. 1E-H). In general, the standard model overestimated both the all IOP components in the blue.

C-modeled amplitudes were linearly correlated with observed values (Fig. 2) and these relationships are expressed by the slope (m) and intercept (b). The observations span a large dynamic range and because correlation coefficients are driven by range and not localized fit, the mean absolute deviation (mad) of retrieved estimates about the linear fit is a robust estimator of goodness-of-fit [Press *et al.*, 1989] (Table 1). C-model-derived $c_p(440)$ and $a_p(440)$ estimates were highly correlated with observed values. C-model-derived estimates of the particle backscattering ratio were not as strongly correlated with observed values but the fit improved when 2 observations associated with dominance by large detrital organic particles are excluded (0.77, 0.001, 0.001 for m, b, mad, respectively). C-model-derived estimates of γ had the weakest correlation, particularly for cases where particles were dominated by extreme concentrations of large ($>20\mu\text{m}$) algal cells. The fit improved significantly without them (1.01, 0.11, 0.10 for m, b, mad, respectively).

Table 1. Comparison of modeled and observed amplitudes, m, b, and mad are the slope, intercept, and mean absolute deviation, respectively. mad for a_p and c_p in units of m^{-1} .

Amplitude	Range	m	b	mad
$c_p(440)$	0.08 - 5.19	0.82	0.26	0.30
$a_p(440)$	0.005 - 1.935	1.02	0.02	0.15
\tilde{b}_{bp}	0.004 - 0.018	0.57	0.003	0.002
γ	0.76 - 0.24	0.68	0.15	0.22

4. Discussion

The importance of obtaining more than chlorophyll from ocean color has contributed to the number of inversion algorithms. Enhanced spectral resolution of remote and in situ radiometers has expanded the capability for retrieving not only more components by inversion, but improved spectral information associated with those components. Regardless, published inversion models have yet to remove the constraints on particle backscattering spectra. In this paper we present an inversion algorithm in which the spectral constraints on the particle backscattering are removed and placed on the particle attenuation coefficient and the backscattering ratio, two parameters with better-established spectral dependences [Voss, 1992; Barnard *et al.*, 1998; Boss *et al.*, 2001b]. The implicit assumption is that the particles responsible for backscattering

are the same particles responsible for attenuation and absorption. Whereas theoretically each IOP is differently dependent on particle size [Stramski and Kiefer, 1991; Boss *et al.*, 2001b], under most conditions it is likely that the same portions of the PSD will dominate all the IOPs, and in the open ocean, where small particles dominate, the IOPs and particles are well correlated. Future field, laboratory and modeling work will provide better constraints for these model assumptions.

Under conditions where all optical properties are dominated by algal cells, we expect the backscattering to vary predictably and in concert with absorption as has been documented for algal cultures. The c-model applied to extreme algal blooms yields these predictable backscattering spectral signatures that cannot be retrieved using a hyperbolic function for backscattering. The standard model did not provide realistic backscattering spectra although it did fit the measured reflectance spectra reasonably well. This was due to variance transference between the particle backscattering and the CDM and CPM absorption basis functions. Transference is a common occurrence in non-linear least squares problems in which exponentially (or hyperbolically) varying basis functions occur in both the numerator and the denominator. Minimization is achieved by balance between the two.

An important achievement of the c-model is the estimate of c_p within 0.23 m^{-1} (average deviation 30%) over two orders of dynamic range. These results might be surprising given the lack of an analytic relationship between R and c (R is dominated by the small proportion of scattering that occurs at large angles while c is dominated by the large proportion of scattering at the smallest angles; [Gordon, 1993]), except that over a large dynamic range there is a first order relationship between backscattering and forward scattering driven by mass concentration. Higher order variations are induced by PSD and the real refractive index, but will be captured by \tilde{b}_{bp} , which is estimated with an average deviation of 17% (maximally 50%).

This inversion does not depend on or constrain the relationship between c_p and b_{bp} , which are differently impacted by particle size and composition; in fact no relationship was observed in this data set. Thus, this model provides a significant improvement compared to empirical estimation of beam attenuation from backscattering coefficients. Our interest in retrieving c_p from ocean color or surface reflectance lies in its strong correlation with particle volume and particulate organic carbon [Spinrad, 1986; Bishop, 1999]. The poorest derived quantity from the c-model is γ , although values were estimable within 0.22. Since γ is related linearly to the Junge slope of the PSD for most oceanic conditions [Volz, 1954; Boss *et al.*, 2001a, b], retrieved values may be sufficient for qualitative estimates of PSD and ecosystem structure.

Extending this model to remotely sensed ocean color data will require sensitivity analyses, incorporating uncertainty due to atmospheric correction and testing on multispectral data. However, the model in its current form is applicable to surface and near surface observations

Acknowledgments. NASA grant NAG5-7872 (CSR) and support from the ONR Environmental Optics Program (EB) funded this work. CSR would like to thank G. Pitcher for the Benguelan field

experiment, M. Sieracki for the Gulf of Maine cruise and S. Etheridge and H. Franklin for valuable assistance.

References

- Barnard, A. H., W. S. Pegau, and J. R. V. Zaneveld, Global relationships in the inherent optical properties of the oceans, *J. Geophys. Res.*, *103*, 24,955-24,968, 1998.
- Bishop, J. K. B., Transmissometer measurement of POC. *Deep-Sea Res.*, *46*, 353-369, 1999.
- Boss, E., W. S. Pegau, W. D. Gardner, J. R. V. Zaneveld, A. H. Barnard, M. S. Twardowski, G. C. Chang, and T. D. Dickey, Spectral particulate attenuation and particle size distribution in the bottom boundary layer of a continental shelf, *J. Geophys. Res.*, *106*, 9509-9516, 2001a.
- Boss E., M. S. Twardowski and S. Herring, The shape of the particulate beam attenuation spectrum and its relation to the size distribution of oceanic particles, *Appl. Opt.*, *40*, 4885-4893, 2001b.
- Bricaud, A., and A. Morel, Light attenuation and scattering by a phytoplanktonic cells: A theoretical modeling, *Appl. Opt.*, *25*, 571-580, 1986.
- Garver, S. A. and D. A. Siegel, Inherent optical property inversion of ocean color spectra and its biogeochemical interpretation. I. Time series from the Sargasso Sea, *J. Geophys. Res.*, *102*, 18,607-18,625, 1997.
- Gordon, H. R., Spectral variations in the volume scattering function at large angles in natural waters, *J. Opt. Soc. Am.*, *64*, 773-775, 1974.
- Gordon, H. R., The sensitivity of radiative transfer to small-angle scattering in the ocean: A quantitative assessment, *Appl. Optics*, *32*, 7505-7511, 1993.
- Gordon, H. R., O. B. Brown, R. H. Evans, J. W. Brown, R. C. Smith, K. S. Baker, and D. K. Clark, A semianalytic radiance model of ocean color, *J. Geophys. Res.*, *93*, 10,909-10,924, 1988.
- Hoge, F. E., and P. E. Lyon, Satellite retrieval of inherent optical properties by linear matrix inversion of oceanic radiance models: An analysis of model and radiance measurement errors, *J. Geophys. Res.*, *101*, 16,631-16,648, 1996.
- Hoge, F. E., and P. E. Lyon, Spectral parameters of inherent optical property models: method for satellite retrieval by matrix inversion of an oceanic radiance model, *Appl. Optics*, *38*, 1657-1662, 1999.
- Lee, Z., K. L. Carder, S. K. Hawes, R. G. Steward, T. G. Peacock, and C. O. Davis, Model for the hyperspectral remote-sensing reflectance, *Appl. Opt.*, *33*, 5721-5732, 1994.
- Morel, A., Diffusion de la lumière par les eaux de mer. Résultats expérimentaux et approach théorique, *Optics of the Sea*, pp. 31.1-31.76, NATO AGARD lecture series No. 61, 1973.
- Morel, A., Optical properties of pure water and pure seawater, in *Optical Aspects of Oceanography* edited by N. G. Jerlov and E. S. Nielsen, pp. 1-24, Academic, 1974.
- Morel, A. and B. Gentili, Diffuse reflectance of oceanic waters. III. Implication of bidirectionality for the remote-sensing problem, *Appl. Opt.*, *35*, 4850-4862, 1996.
- Morel, A., D. Antoine and B. Gentili, Bidirectional reflectance of oceanic waters: Accounting for Raman emission and varying particle phase function, *Appl. Opt.*, *41*, 6289-6306, 2002.
- Pegau, W. S., D. Gray, and J. R. V. Zaneveld. 1997. Absorption and attenuation of visible and near-infrared light in water: Dependence on temperature and salinity. *Appl. Opt.* *36*: 6035-6046.
- Petzold, T. J., Volume scattering functions for selected ocean waters, No. Ref. 72-78, 79pp. Scripps Inst. Oceanogr, 1972.
- Pope, R. M. and E. S. Fry, Absorption spectrum (380-700 nm) of pure water. II. Integrating cavity measurements. *Appl. Opt.*, *36*, 8710-8723, 1997.
- Press, W. H., B. P. Flannery, S. A. Teukolsky, and W. T. Vetterling, *Numerical Recipes, the Art of Scientific Computing*. Cambridge, 1989.

- Roesler, C. S., Theoretical and experimental approaches to improve the accuracy of particulate absorption coefficients from the Quantitative Filter Technique, *Limnol. Oceanogr.*, *43*, 1649-1660, 1998.
- Roesler, C. S., S. M. Etheridge, and G. P. Pitcher, Application of an ocean color algal taxa detection model to the red tide communities of the southern Benguela, Submitted, Proc. 10th Intl. Conf. Harm Algal Blooms, in press 2003.
- Roesler, C. S. and M. J. Perry, In situ phytoplankton absorption, fluorescence emission, and particulate backscattering spectra determined from reflectance, *J. Geophys. Res.*, *100*, 13,279-13,294, 1995.
- Roesler, C. S., M. J. Perry, and K. L. Carder, Modeling *in situ* phytoplankton absorption from total absorption spectra, *Limnol. Oceanogr.*, *34*, 1512-1525, 1989.
- Spinrad, R. W., An optical study of the water masses of the Gulf of Maine, *J. Geophys. Res.*, *91*, 1007-1018, 1986.
- Stramski, D. and D. A. Kiefer, Light scattering by microorganisms in the open ocean, *Prog. Oceanogr.*, *28*, 343-383, 1991.
- Stramski, D. and C. D. Mobley, Effects of microbial particles on oceanic optics: a database of single-particle optical properties, *Limnol. Oceanogr.*, *42*, 538-549, 1997.
- Twardowski M., E. Boss, J. B. Macdonald, W. S. Pegau, A. H. Barnard, and J. R. V. Zaneveld, A model for estimating bulk refractive index from the optical backscattering ratio and the implications for understanding particle composition in case I and case II waters, *J. Geophys. Res.*, *106*, 14,129-14,142, 2001.
- Ulloa O., S. Sathyendranath and T. Platt, Effect of the particle-size distribution on the backscattering ratio in seawater, *Appl. Opt.*, *33*, 7070-7077, 1994.
- van de Hulst, H. C., *Light Scattering by Small Particles*. Wiley, 1957.
- Volz, F. 1954. Die optik und meteorologie der atmospharischen Trubung. Ber. Dtsch. Wetterdiensstes 2: 3-47.
- Voss, K. J., A spectral model of the beam attenuation coefficient in the ocean and coastal areas, *Limnol. Oceanogr.*, *37*, 501-509, 1992.
- Zaneveld, J. R. V., and J. C. Kitchen, The variation in the inherent optical properties of phytoplankton near an absorption peak as determined by various models of cell structure, *J. Geophys. Res.*, *100*, 13,309-13,320, 1995. Belt, E. S., Post-Acadian rifts and related facies, eastern Canada, in *Studies in Appalachian Geology*, edited by E. Zen et al., pp. 95-113, John Wiley, New York, 1968.

Mailing addresses: E. Boss, School of Marine Sciences, 5741 Libby Hall, University of Maine, Orono, ME 04469, C. Roesler, Bigelow Laboratory for Ocean Sciences, POBox 475, West Boothbay Harbor, ME 04575

(Received October 6, 2000; revised February 27, 2001; accepted March 31, 2001.)

AGU Copyright:

Copyright 2001 by the American Geophysical Union.

Paper number 2001GL012345.
0094-8276/01/2001GL012345\$05.00

Public Domain Copyright:

3.09 This paper is not subject to U.S. copyright. Published in 2001 by the American Geophysical Union.

Paper number 2001GL012345.

Crown Copyright:

Published in 2001 by the American Geophysical Union.

Paper number 2001GL012345.

****Provide running head (45 character max for short title):**

**ROESLER AND BOSS: BEAM ATTENUATION FROM
OCEAN COLOR**




Figure 1. Observed and modeled parameters for two optically diverse regimes: spectral reflectance (A, irradiance, and B, radiance, reflectance), particulate attenuation (C, D), particulate absorption (E, F) and CDM absorption (G, H). Observations were made in the gyre waters off the Oregon coast (left panels, data and absorption methods from *Roesler and Perry* [1995]), and Benguelian coastal waters off South Africa (right panels). Chlorophyll values were 0.07 and 131.65 $\mu\text{g/L}$, respectively. Symbols: observed (black or symbols), c-model (dark stipple), standard model (light stipple).




Figure 2. Relationship between measured and modeled optical properties. Particulate attenuation (A) and absorption coefficient (B) at 440nm, particulate backscattering ratio at 650nm (C; 532 nm for gyre and GoM), and slope of the particulate attenuation spectrum (γ , D).